

Vitamin D for prevention of respiratory tract infections: A systematic review and meta-analysis

Jaykaran Charan, Jagdish P. Goyal¹, Deepak Saxena², Preeti Yadav

Departments of Pharmacology, ¹P. D. U. Govt. Medical College, Rajkot, ²Department of Epidemiology, Indian Institute of Public Health, Ahmedabad, Gujarat, India

ABSTRACT

Objectives: To explore the effect of vitamin D supplementation in prevention of respiratory tract infections on the basis of published clinical trials. **Materials and Methods:** Clinical trials were searched from various electronic databases. Five clinical trials were suitable for inclusion. Outcome was events of respiratory tract infections in vitamin D group and placebo group. Data was reported as odds ratio with 95% confidence interval. Both random and fixed model was used for analysis. Analysis was done with the help of Comprehensive meta-analysis software 2. **Results:** Events of respiratory tract infections were significantly lower in vitamin D group as compared to control group [Odds ratio = 0.582 (0.417 – 0.812) $P=0.001$] according to random model. Results were similar in fixed model. On separate analysis of clinical trials dealing with groups of children and adults, beneficial effect of vitamin D was observed in both, according to fixed model [Odds ratio = 0.579 (0.416 – 0.805), $P=0.001$ and Odd ratio = 0.653 (0.472 – 0.9040, $P=0.010$ respectively]. On using random model beneficial effect persisted in children's group but became nonsignificant in adults group [Odds ratio = 0.579 (0.416 – 0.805), $P=0.001$ and Odd ratio = 0.544 (0.278 – 1.063) $P=0.075$ respectively]. **Conclusion:** Vitamin D supplementation decreases the events related to respiratory tract infections. There is need of more well conducted clinical trials to reach to a certain conclusion.

Key words: Influenza, pneumonia, prevention, respiratory tract infections, vitamin D

INTRODUCTION

Acute respiratory infection (ARI) is widespread not only in the pediatric age group but also in adults. Respiratory tract infections are the major cause of mortality and morbidity.^[1]

There are limited options available for treatment and prevention of ARI. Antibiotics are the mainstay of treatment

for pneumonia. In the developing countries, zinc and vitamin A supplementation has been studied for the prevention and treatment of ARI. It has been shown that supplements of these micronutrients reduce the ARI rates though evidences are conflicting.^[2-4] Vitamin C (ascorbic acid) has also been used for the prevention and treatment of ARI despite the lack of convincing evidence of benefit.^[5]

Recent research suggested that vitamin D also has a potential role in the prevention of ARI by increasing immunity. In some observational studies, it was observed that low vitamin D level in blood is associated with the increased incidence of respiratory tract infections.^[6] There is a need to refine our knowledge about the functions of vitamin D in the prevention of respiratory tract infections in the light of clinical trials which were done to explore this phenomenon. Hence, we conducted

Access this article online	
Quick Response Code:	Website: www.jpharmacol.com
	DOI: 10.4103/0976-500X.103685

Address for correspondence:

Jaykaran Charan, Department of Pharmacology, Govt. Medical College, Surat, Gujarat, India.
E-mail: drjaykaran@yahoo.co.in

this systematic review and meta-analysis to assess the role of vitamin D supplementation in prevention of ARI.

MATERIALS AND METHODS

Search strategy of clinical trials

This systematic review and meta-analysis was done according to the instructions given in “Cochrane Handbook for Systematic Reviews of Interventions”.^[7] Clinical trials were collected from the electronic databases like PubMed, Cochrane Clinical Trial Register, and Google Scholar. References mentioned in full text articles and reviews were also explored to collect the relevant clinical trials. The clinical trials which are not available as free full text were collected from the corresponding authors of these trials through email. In PubMed, various MeSH terms like “Vitamin D”, “Respiratory Tract Infections”, “Influenza”, “Pneumonia”, “Common Cold”, etc., were used in various combinations to retrieve the clinical trials. For PubMed, clinical trials published from 1966 to present were considered. Only those clinical trials which are published in English language were collected. Summary of various words used for searching the literature is given in appendix 1. Last search was done on 23 November 2011. Both first and second authors [J.K. and J.P.G.] did the search together from databases. Potential clinical trials were downloaded in hard copies. The clinical trials which were not available online in full text were collected from the corresponding author through email. These clinical trials were assessed on the basis of inclusion and exclusion criteria mentioned in the pretested proforma independently by both authors. Wherever there was discrepancy, the decision of the third author was considered final.

Inclusion and exclusion criteria for clinical trials

Only randomized placebo controlled clinical trials exploring the effect of vitamin D supplementation on prevention of respiratory tract infections like influenza, pneumonia, common cold, etc., were included in the analysis. Only those clinical trials where results were given in categorical variables were included. Non randomized, clinical trials without controls and clinical trials showing results in continuous variables were not included. Clinical trials in which vitamin D was used in combination with other intervention/s were not included in the final analysis. Process of selection of various RCTs for this study is given in the form of preferred reporting items for systematic review and meta-analysis (PRISMA) chart according to PRISMA statement [Figure 1].^[8]

Outcome

Outcome of interest was the event of respiratory tract infection in the case and control groups. In all clinical trials placebo was used in the control group. The software used in this analysis (Comprehensive meta-analysis software 2) converted these events into odds ratio and calculated 95% confidence interval around odds ratio.

Statistics

The statistical analysis was performed by CMA (Comprehensive meta-analysis) Version 2 software and the effect was estimated by calculating the odds ratio (OR) and the 95% confidence interval (CI). No data transformation was done. Events in each group were combined for meta-analysis. Results were given in both random as well as fixed model. Forest plot values were given according to random model and test for heterogeneity (τ^2) was done.

RESULTS

On the basis of inclusion and exclusion criteria, we found 5 clinical trials.^[9-14] Dose of vitamin D used in these clinical trials were ranged from 400 IU/day to 2000 IU/day. In one clinical trial single parenteral dose of vitamin D was given (100000 IU). [Table 1] In four clinical trials, events were recorded in the form of number of participants suffered from respiratory tract infections during the trial period. In one clinical trial events were recorded in the form of total number of times participants reported respiratory tract infection in the questionnaire given to them (four times) in trial period. Levels of evidence of these clinical trials were high or moderate. Level of evidence was assessed according to “The Grades of Recommendation, Assessment, Development and Evaluation Working Group” (GRADE Working group).^[15,16] Information related to the quality of evidence and biases in these clinical trials are mentioned in Table 2.

Overall events of respiratory tract infection in vitamin D group were 21.7% (205/943) whereas in placebo group it was 30.1% (279/925). There was significant reduction in incidence of respiratory tract infection in vitamin D group as compared to placebo group [Odds ratio = 0.582 (0.417 – 0.812) $Z = -3.185$, P value = 0.001, $\tau^2 = 0.064$] according to the random model [Figure 2]. The values were similar when fixed model was used [Odds ratio = 0.615 (0.488 – 0.776) $Z = -4.108$, P value = 0.000].

There were two clinical trials which were dealing with pediatric age group. These clinical trials were analyzed separately and it was observed that in this age group also vitamin D




Figure 1: PRISMA Chart

supplementation reduces the rate of respiratory tract infections significantly [Odds ratio = 0.579 (0.416 – 0.805) $Z = -3.246$, P value = 0.001 test for heterogeneity $t^2 = 0.000$] by random model [Figure 3]. After using the fixed model, values were same [Odds ratio = 0.579 (0.416 – 0.805) $Z = -3.246$, P value = 0.001].

Three clinical trials were dealing with adult population. In these clinical trials, odd ratio was in favor of vitamin D group but it was not significant. [Odd ratio = 0.544 (0.278 – 1.063) $Z = -1.783$, P value = 0.075, test for heterogeneity $t^2 = 0.247$] according to random model. [Figure 3] After analyzing on the basis of fixed model, values were significantly in favor of vitamin D [Odd ratio = 0.653 (0.472 – 0.904) $Z = -2.569$, P value = 0.010].

Publication bias (through funnel plot) was not checked because of small numbers of clinical trials.^[17]

DISCUSSION

This systematic review and meta-analysis was done to explore the role of vitamin D supplementation in prevention of respiratory tract infections. According to this systematic review and meta-analysis vitamin D significantly reduces the respiratory tract infection related events as compared to placebo. Beneficial effect of vitamin D was observed in children as well as adults according to fixed model. Beneficial effect of vitamin D was observed only in children according to random model. This assessment is based on only five clinical trials as less clinical trials are done for the exploration of the same.

Various reasons are given for the protective action of vitamin D in prevention of respiratory tract infections. It is believed that vitamin D increases the production of natural antibodies.^[18]

Vitamin D is also known to strengthen the immunity by inducing monocyte differentiation and inhibiting lymphocyte proliferation.^[19] It also postulated that vitamin D enhances the phagocytic activity of macrophages.^[18]

This systematic review and meta-analysis has some limitations. In clinical trials dealing with children there was significant protective effect of vitamin D in our analysis but there was




Figure 2: Meta-analysis of trials showing effect of vitamin D supplementation in prevention of respiratory tract infections. Effect of vitamin D supplementation in prevention of respiratory tract infections. Random effect model (DerSimonian laird); Odds ratio = 0.582 (0.417 – 0.812) $Z = -3.185$, P value = 0.001, test for heterogeneity $t^2 = 0.064$




Figure 3: Meta-analysis of trials showing effect of vitamin D supplementation in prevention of respiratory tract infections in pediatric population. Effect of vitamin D supplementation in prevention of respiratory tract infections in pediatric population. Random effect model (DerSimonian laird); Odds ratio = 0.579 (0.416 – 0.805) $Z = -3.246$ P Value = 0.001 Test for heterogeneity $t^2 = 0.000$

Table 1: Characteristics of clinical trials included in the meta-analysis

Name	Dose	Duration	Event in drug	Event in control	Population
Ilkka Laaksi <i>et al.</i> , 2010	400 IU/Day	6 months	39/80	50/80	18-28 year men
Mitsuyoshi Urashima <i>et al.</i> , 2010	1200 IU/Day	4 months	18/167	31/167	6-15 year children
M. LI -NG <i>et al.</i> , 2009	1200 IU/Day	3 months	48/388	50/363	18-80 year adults
Semira Manaseki-Holland <i>et al.</i> , 2010	100000 IU once	3 months	92/204	122/211	1-3 year children
Aloia JF, <i>et al.</i> , 2007	2000 IU/Day	3 year	8/104	26/104	Post menopausal women

Table 2: Levels of evidence and methodological bias in clinical trials included in the meta-analysis

	Ilkka Laaksi <i>et al.</i> , 2010	Mitsuyoshi Urashima <i>et al.</i> , 2010	M. LI -NG <i>et al.</i> , 2009	Semira Manaseki-Holland <i>et al.</i> , 2010	Aloia JF, <i>et al.</i> , 2007
Adequate sequence generation?	Yes	Yes	Yes	Yes	Yes
Allocation concealment?	Yes	Yes	Yes	Yes	No
Blinding?	Yes	Yes	Yes	Yes	Yes
Incomplete outcome data?	No	Yes	Yes	No	Yes
Selective reporting?	No	No	No	No	Unclear
Free of other bias?	Yes	Unclear	Unclear	No	Unclear
Grade of evidence	High	Moderate	Moderate	High	Poor




Figure 4: Meta-analysis of trials showing effect of vitamin D supplementation in prevention of respiratory tract infections in adult population. Effect of vitamin D supplementation in prevention of respiratory tract infections in adult population. Random effect model (DerSimonian laird); Odds ratio = 0.544 (0.278 – 1.063) Z = - 1.783 P value = 0.075 test for heterogeneity $t^2 = 0.247$

significant heterogeneity [Test for heterogeneity $t^2 = 0.000$]. But same odds ratio was observed in random as well as fixed model. This heterogeneity may be there because of some publication bias, very less numbers of clinical trials, different doses of vitamin D and heterogeneity of participants of clinical trials. We could not check publication bias because interpretation of funnel plot is not possible when clinical trials are less than 10.^[17] Because of insufficient reporting we could not conduct the analysis of adverse effects. We could not compare our results with other meta-analyses, as not a single meta-analysis exploring this phenomenon is existed in electronic databases.

On the basis of this study, we can conclude that vitamin D is useful in prevention of respiratory tract infections. But looking at the availability of only five clinical trials there is need of conduction of more clinical trials so that more valid conclusion can be reached.

Appendix 1

Summary of words used in Pubmed search

- Vitamin D/
- Cholecalciferol/
- Cholecalciferol [Text word]
- Ergocalciferol/
- Ergocalciferol [Text word]
- Vitamin D [Text word]
- Respiratory tract infections/
- Respiratory tract infections [Text word]
- Upper respiratory tract infections/
- Upper respiratory tract infections [Text word]
- Respiratory Tract Infections/prevention and control
- Influenza, Human/
- Influenza [Text word]
- Pneumonia/
- Pneumonia [Text word]
- Common cold/
- Common cold [Text word]
- Randomized controlled trials [Publication type]
- Controlled clinical trials [Publication type]

REFERENCES

1. World Health Organization: The global burden of disease: 2004 update. Available from: http://www.who.int/healthinfo/global_burden_disease/GBD_report_2004update_full.pdf [Last accessed on 2011 Nov 18].
2. International Institute for Population Sciences (IIPS) and Macro International. National Family Health Survey(NFHS-3), 2005-06: India. Mumbai: IIPS; 2008.
3. Acharya D, Prasanna KS, Nair S, Rao RS. Acute respiratory infections in children: A community based longitudinal study in south India. Indian J Public Health 2003;47:7-13.
4. Yamey G. Zinc supplementation prevents diarrhoea and pneumonia. BMJ 1999;319:1521-2.
5. Bhutta ZA, Black RE, Brown KH, Gardner JM, Gore S, Hidayat A, *et al.* Prevention of diarrhea and pneumonia by zinc supplementation in developing countries: Pooled analysis of randomized controlled trials. Zinc Investigators' Collaborative Group. J Pediatr 1999;135:689-97.
6. Wayse V, Yousafzai A, Mogale K, Filteau S. Association of subclinical vitamin D deficiency with severe acute lower respiratory infection in Indian children under 5 years. Eur J Clin Nutr 2004;58:563-7.
7. Higgins JP, Green S, editors. Cochrane Handbook for Systematic Reviews of Interventions. Chichester (UK): John Wiley and Sons; 2008.
8. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, *et al.* The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: Explanation and elaboration. PLoS Med 2009;6:e1000100.
9. Laaksi I, Ruohola JP, Mattila V, Auvinen A, Ylikomi T, Pihlajamäki H. Vitamin D Supplementation for the Prevention of Acute Respiratory Tract Infection: A Randomized, Double-Blinded Trial among Young Finnish Men. J Infect Dis 2010;202:809-14.
10. Urashima M, Segawa T, Okazaki M, Kurihara M, Wada Y, Ida H. Randomized trial of vitamin D supplementation to prevent seasonal influenza A in schoolchildren. Am J Clin Nutr 2010;91:1255-60.
11. Holland SM, Qader G, Masher MI, Bruce J, Mughal MZ, Chandramohan D, *et al.* Effects of vitamin D supplementation to children diagnosed with pneumonia in Kabul: A randomised controlled trial. Trop Med Int Health 2010;15:1148-55.
12. Li-NG M, Aloia JF, Pollack S, Cunha BA, Mikhail M, Yeh J, *et al.* A randomized controlled trial of vitamin D3 supplementation for the prevention of symptomatic upper respiratory tract infections. Epidemiol Infect 2009;137:1396-404.
13. Aloia JF, Talwar SA, Pollack S, Yeh J. A randomized controlled trial of vitamin D3 supplementation in African American women. Arch Intern Med 2005;165:1618-23.
14. Aloia J, Li-NG M. Re: Epidemic influenza and vitamin D. Epidemiol Infect 2007;135:1095-6.
15. GRADE Working Group. Grading quality of evidence and strength of recommendations. BMJ 2004;328:1490-4.
16. Schunemann HJ, Oxman AD, Vist GE, Higgins JP, Deeks JJ, Glasziou P, *et al.* Chapter 12: Interpreting results and drawing conclusions. In: Higgins JP, Green S, editors. Cochrane Handbook for Systematic Reviews of Interventions. Chichester (UK): John Wiley and Sons; 2008.
17. Sterne JAC, Egger M, Moher D, editors. Chapter 10: Addressing reporting biases. In: Higgins JPT, Green S (editors), Cochrane Handbook for Systematic Reviews of Interventions. Chichester (UK): John Wiley and Sons; 2008.
18. Raloff J. The antibiotic vitamin. Deficiency in vitamin D may predispose people to infection. Science 2006;170:312.
19. Liu PT, Stenger S, Li H, Wenzel L, Tan BH, Krutzik SR, *et al.* Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. Science 2006;311:1770-3.

How to cite this article: Charan J, Goyal JP, Saxena D, Yadav P. Vitamin D for prevention of respiratory tract infections: A systematic review and meta-analysis. J Pharmacol Pharmacother 2012;3:300-3.

Source of Support: Nil, **Conflict of Interest:** None declared.